

Order Plecoptera



	<p>Family Perlidae Golden/common stonefly (5-20 mm)</p> <p>Usually large and strikingly patterned; finely branched gills present on all thoracic segments; wing pads diverge slightly from the midline. Occur in fast areas of streams in rivers on rocks and submerged logs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>		<p>Family Perlidae Golden/common stonefly (5-20 mm)</p> <p>Usually large and strikingly patterned; finely branched gills present on all thoracic segments; wing pads diverge slightly from the midline. Occur in fast areas of streams in rivers on rocks and submerged logs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>
---	--	---	--



	<p>Family Perlidae Golden/common stonefly (5-20 mm)</p> <p>Usually large and strikingly patterned; finely branched gills present on all thoracic segments; wing pads diverge slightly from the midline. Occur in fast areas of streams in rivers on rocks and submerged logs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>		<p>Family Perlidae Golden/common stonefly (5-20 mm)</p> <p>Usually large and strikingly patterned; finely branched gills present on all thoracic segments; wing pads diverge slightly from the midline. Occur in fast areas of streams in rivers on rocks and submerged logs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>
--	--	--	--

Order Plecoptera

	<p>Family Capniidae Small winter stonefly (5-20 mm)</p> <p>Slender elongated body; front of thorax slightly wider than the abdomen; wing pads not divergent from the midline; abdominal segments separated by a membranous fold. Occur in small and medium sized rivers and streams with gavel-cobble substrates and fast flows.</p> <p>Tolerance: Low Behavior: Clinger (crawler) Feeding: Shredder</p>		<p>Family Capniidae Small winter stonefly (5-20 mm)</p> <p>Slender elongated body; front of thorax slightly wider than the abdomen; wing pads not divergent from the midline; abdominal segments separated by a membranous fold. Occur in small and medium sized rivers and streams with gavel-cobble substrates and fast flows.</p> <p>Tolerance: Low Behavior: Clinger (crawler) Feeding: Shredder</p>
	<p>Family Capniidae Small winter stonefly (5-20 mm)</p> <p>Slender elongated body; front of thorax slightly wider than the abdomen; wing pads not divergent from the midline; abdominal segments separated by a membranous fold. Occur in small and medium sized rivers and streams with gavel-cobble substrates and fast flows.</p> <p>Tolerance: Low Behavior: Clinger (crawler) Feeding: Shredder</p>		<p>Family Capniidae Small winter stonefly (5-20 mm)</p> <p>Slender elongated body; front of thorax slightly wider than the abdomen; wing pads not divergent from the midline; abdominal segments separated by a membranous fold. Occur in small and medium sized rivers and streams with gavel-cobble substrates and fast flows.</p> <p>Tolerance: Low Behavior: Clinger (crawler) Feeding: Shredder</p>

Order Plecoptera

	<p>Family Perlodidae Patterned stonefly (10-50 mm)</p> <p>Strikingly patterned and colored similar in appearance to Perlidae; hind wing pads divergent; no gills on the thoracic segments. Occur in fast flowing streams and rivers under rocks, submerged logs and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>		<p>Family Perlodidae Patterned stonefly (10-50 mm)</p> <p>Strikingly patterned and colored similar in appearance to Perlidae; hind wing pads divergent; no gills on the thoracic segments. Occur in fast flowing streams and rivers under rocks, submerged logs and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>
---	--	---	--



	<p>Family Perlodidae Patterned stonefly (10-50 mm)</p> <p>Strikingly patterned and colored similar in appearance to Perlidae; hind wing pads divergent; no gills on the thoracic segments. Occur in fast flowing streams and rivers under rocks, submerged logs and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>		<p>Family Perlodidae Patterned stonefly (10-50 mm)</p> <p>Strikingly patterned and colored similar in appearance to Perlidae; hind wing pads divergent; no gills on the thoracic segments. Occur in fast flowing streams and rivers under rocks, submerged logs and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Predator</p>
--	--	--	--


Order Plecoptera

	<p>Family Peltoperlidae Roach-like stonefly ^(5-15 mm)</p> <p>Small stout body; rear divergent wing pads; thoracic segments are oval or triangular shaped and cover much of the upper body; some have fine gills on the front legs. Occur in fast flowing water on roots, leaf packs and rocks.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>		<p>Family Peltoperlidae Roach-like stonefly ^(5-15 mm)</p> <p>Small stout body; rear divergent wing pads; thoracic segments are oval or triangular shaped and cover much of the upper body; some have fine gills on the front legs. Occur in fast flowing water on roots, leaf packs and rocks.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
---	--	---	--

	<p>Family Peltoperlidae Roach-like stonefly ^(5-15 mm)</p> <p>Small stout body; rear divergent wing pads; thoracic segments are oval or triangular shaped and cover much of the upper body; some have fine gills on the front legs. Occur in fast flowing water on roots, leaf packs and rocks.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>		<p>Family Peltoperlidae Roach-like stonefly ^(5-15 mm)</p> <p>Small stout body; rear divergent wing pads; thoracic segments are oval or triangular shaped and cover much of the upper body; some have fine gills on the front legs. Occur in fast flowing water on roots, leaf packs and rocks.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
--	--	--	--

Order Plecoptera

	<p>Family Pteronarcyidae Giant stonefly ^(15-70 mm)</p> <p>Very large and usually dark brown in color; finely branched gills on all thoracic segments plus the first two abdominal segments. Occur in swift flowing streams in rivers, mostly in the areas of snags and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>		<p>Family Pteronarcyidae Giant stonefly ^(15-70 mm)</p> <p>Very large and usually dark brown in color; finely branched gills on all thoracic segments plus the first two abdominal segments. Occur in swift flowing streams in rivers, mostly in the areas of snags and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
---	--	---	--

	<p>Family Pteronarcyidae Giant stonefly ^(15-70 mm)</p> <p>Very large and usually dark brown in color; finely branched gills on all thoracic segments plus the first two abdominal segments. Occur in swift flowing streams in rivers, mostly in the areas of snags and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>		<p>Family Pteronarcyidae Giant stonefly ^(15-70 mm)</p> <p>Very large and usually dark brown in color; finely branched gills on all thoracic segments plus the first two abdominal segments. Occur in swift flowing streams in rivers, mostly in the areas of snags and leaf packs.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
--	--	--	--

Order Plecoptera

 	<p>Family Nemouridae Little brown stonefly (5-20 mm)</p> <p>Very small, often hairy appearance; wing pads diverge greatly from the midline; hind legs as long as the abdomen; gills often present between the head and thorax. Common in small coldwater streams, but also occur in larger fast flowing rivers.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>	 	<p>Family Nemouridae Little brown stonefly (5-20 mm)</p> <p>Very small, often hairy appearance; wing pads diverge greatly from the midline; hind legs as long as the abdomen; gills often present between the head and thorax. Common in small coldwater streams, but also occur in larger fast flowing rivers.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
--	--	--	--

 	<p>Family Nemouridae Little brown stonefly (5-20 mm)</p> <p>Very small, often hairy appearance; wing pads diverge greatly from the midline; hind legs as long as the abdomen; gills often present between the head and thorax. Common in small coldwater streams, but also occur in larger fast flowing rivers.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>	 	<p>Family Nemouridae Little brown stonefly (5-20 mm)</p> <p>Very small, often hairy appearance; wing pads diverge greatly from the midline; hind legs as long as the abdomen; gills often present between the head and thorax. Common in small coldwater streams, but also occur in larger fast flowing rivers.</p> <p>Tolerance: Low Behavior: Clinger Feeding: Shredder</p>
---	--	---	--