


Phylum Platyhelminthes

	<p>Class Turbellaria Flatworms (4-450 mm)</p> <p>Soft-elongated body flattened from top to bottom; no individual segments; mouth usually on the bottom side positioned about one-fifth to the length of the body; sides of the body constricted towards the front forming a head that is often somewhat triangular shaped; two eyespots situated on top of the head gives the animal a cross-eyed appearance.</p> <p>Tolerance: High Behavior: Clinger (swimmer) Feeding: Predator (collector)</p>		<p>Class Turbellaria Flatworms (4-450 mm)</p> <p>Soft-elongated body flattened from top to bottom; no individual segments; mouth usually on the bottom side positioned about one-fifth to the length of the body; sides of the body constricted towards the front forming a head that is often somewhat triangular shaped; two eyespots situated on top of the head gives the animal a cross-eyed appearance.</p> <p>Tolerance: High Behavior: Clinger (swimmer) Feeding: Predator (collector)</p>
	<p>Class Turbellaria Flatworms (4-450 mm)</p> <p>Soft-elongated body flattened from top to bottom; no individual segments; mouth usually on the bottom side positioned about one-fifth to the length of the body; sides of the body constricted towards the front forming a head that is often somewhat triangular shaped; two eyespots situated on top of the head gives the animal a cross-eyed appearance.</p> <p>Tolerance: High Behavior: Clinger (swimmer) Feeding: Predator (collector)</p>		<p>Class Turbellaria Flatworms (4-450 mm)</p> <p>Soft-elongated body flattened from top to bottom; no individual segments; mouth usually on the bottom side positioned about one-fifth to the length of the body; sides of the body constricted towards the front forming a head that is often somewhat triangular shaped; two eyespots situated on top of the head gives the animal a cross-eyed appearance.</p> <p>Tolerance: High Behavior: Clinger (swimmer) Feeding: Predator (collector)</p>